

**IGIV and General Immune Disorders
Enrollment Form**

SUPERIOR BIOLOGICS
 Fax Referral To: 914-747-1170
 Phone: 855-747-1150



Date: _____

PATIENT INFORMATION

Patient Name: _____
 Address: _____
 City, State, Zip: _____
 Home Phone: _____
 Cell Phone: _____
 Alternate Phone: _____
 Date of Birth: _____ Gender: M F

PRESCRIBER INFORMATION

Prescriber Name: _____
 Address: _____
 City, State, Zip: _____
 Phone: _____
 Fax: _____
 DEA#: _____ NPI#: _____
 Contact Person: _____

INSURANCE INFORMATION (Please attach the front and back of insurance and prescription drug card)

Primary Insurance: _____ ID#: _____ Group: _____
 Secondary Insurance: _____ ID#: _____ Group: _____
 Prescription Card: _____ ID#: _____ BIN: _____ PCN: _____ Group: _____

DIAGNOSIS (ICD-10)

Neurological

- G61.81 Chronic Inflammatory Demyelinating Polyneuropathy (CIDP)
- G61.82 Multifocal Motor Neuropathy (MMN)
- G61.0 Guillain-Barre G25.82 Stiff-Person Syndrome
- G35 Multiple Sclerosis M33.20 Polymyositis
- G70.01 Myasthenia Gravis w/Exacerbation
- Other: _____

Immunological

- Primary Immune Deficiency – **Please specify ICD-10 Code:** _____
- D80.9 Deficiency of Humoral Immunity
- D83.9 Common Variable Immunodeficiency
- D89.9 Immune Mechanism Disorder D81.9 Immune Deficiency NOS
- D69.3 Idiopathic Thrombocytopenia D80.1 Hypogammaglobulinemia
- Other: _____

CLINICAL INFORMATION (Please attach all clinical information, lab results, and other medical history documents)

Patient Weight: _____ Kg/Lbs Height: _____ Inches/CM Allergies: _____
 Has patient previously received IVIG Yes No Line Access: PIV PICC PORT Needs by Date: _____

Medication	Dose	Directions
<p style="text-align: center;">Intravenous</p> <input type="checkbox"/> Gammagard® Liq. 10% <input type="checkbox"/> Privigen® 10% <input type="checkbox"/> Gamunex-C® 10% <input type="checkbox"/> Octagam® 5% <input type="checkbox"/> Gammaked® 10% <input type="checkbox"/> Octagam® 10% <input type="checkbox"/> Bivigam® <input type="checkbox"/> Gammagard® S/D <input type="checkbox"/> Other: _____ <input type="checkbox"/> Panzyga® 10%	<p>_____ grams OR _____ gram(s) per kg (Pharmacy to round to nearest vial size) Infuse total dose OVER _____ day(s); Every _____ week(s) for: <input type="checkbox"/> 1 month <input type="checkbox"/> 3 months <input type="checkbox"/> 6 months <input type="checkbox"/> 12 months <input type="checkbox"/> Other _____</p>	<p>Infuse total dose of Immunoglobulin intravenously based on manufacturer recommend infusion rate as tolerated. Infuse via: <input type="checkbox"/> Infusion Pump <input type="checkbox"/> Gravity</p>

Medication	Dose	Directions
<p style="text-align: center;">Subcutaneous</p> <input type="checkbox"/> Gammagard® Liq. 10% <input type="checkbox"/> Xembify® 20% <input type="checkbox"/> Gamunex-C® 10% <input type="checkbox"/> Cutaquig® 16.5% <input type="checkbox"/> Gammaked® 10% <input type="checkbox"/> Hizentra® 20% <input type="checkbox"/> HyQvia® 10%	<p>_____ grams OR _____ gram(s) per kg (Pharmacy to round to nearest vial size) Infuse total dose OVER _____ day(s); Every _____ week(s) for: <input type="checkbox"/> 1 month <input type="checkbox"/> 3 months <input type="checkbox"/> 6 months <input type="checkbox"/> 12 months <input type="checkbox"/> Other _____</p>	<p>Infuse total dose of Immunoglobulin subcutaneously in one or more infusion sites via infusion pump based on manufacturer recommend infusion rate as tolerated. Other: _____</p>

Premedication to be given 30 minutes prior to infusion:
 Diphenhydramine 25-50 mg po – 25mg #2 per dose
 Diphenhydramine 25-50 Slow IV-Push – 50mg vial #1 per dose
 Acetaminophen 325-650 mg po – 325mg #2 per dose
 Ketorolac 30mg Slow IV-Push – 30mg/ml vial #1 per dose
 LMX-4 Cream – apply topically to insertion site as needed. #1 tube
 Other: _____

IV Access Flush Order / EpiPen® Order: (Infusion supplies per pharmacy protocol)
 NaCl 0.9% 5-10ml IV before and after infusion
 Heparin 10 units/ml 3-5ml IV after infusion for peripheral access and PRN
 Heparin 100 units/ml 3-5ml IV after infusion for central IV access and PRN
 All infusion supplies necessary to administer the medication
 EpiPen® 0.3mg auto-injector for severe anaphylactic reaction for patient weighing ≥ 30kg. EpiPen Jr. @ 0.15mg for patients weighing under 30kg

By signing below, I certify that above therapy is medically necessary. **Prescriber's Signature (SIGN BELOW)**

 Dispense as Written Date Substitution Allowed Date

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately.