Crohn's and Ulcerative Colitis		SUPER	SUPERIOR BIOLOGICS								
Referral Form		Fax Referral To: 914-747-11		o: 914-747-1170	SUPE	RIC	R				
Phone: 855 747 1150											
Patient Information Prescriber Information											
Please complete the following or send patient demographic she											
Patient Name:											
				City, State, Zip:							
	p:			Phone:							
-	۲۰			DEA: NPI#:							
				Contact Person:							
DOB:	Gender:										
B0B			nca	nformation							
Drimon (Inc.)	ranaal				Croup						
	surance: Card: ID#:										
	Card:ID#: dical Information (Section mus	the completed			Group: (Attach separate sheet if	ineede	d)				
	zation Insurance Number:	t be completed	to pro	cess prescription)	(Allach separate sneeth	neede	a)				
	Please include diagnosis name	with ICD-10 c	ode	Therapy Details:	ew	estart					
	<u> </u>										
🗆 K50.00 Cr	ohn's disease of small intestines wi	thout complicat	tions	Weightk	/lbs Height		_cm/in				
□ K50.8Cro	ohn's disease of both intestines wit	houtcomplicat	ions	Allergies							
🗆 K50.10 Cr	ohn's disease of large intestines wi	thout complicat	ions	Lab Data							
🗆 K50.00 Cr	rohn's disease, unspecified, withc	out complication	ns	Prior Therapies							
□ K20.0 Eosinophilic Esophagitis				Concomitant Medications							
□ Other diagnosis: ICD-10 code				Additional Comments							
Description	Date of Description	า		Injection Training Requi	red? □Yes □No						
Has a TB test	t been performed?	Yes 🗆	No								
Does the Pat	ient have an active infection? \Box	Yes 🗆 N	lo								
Start Date	Review Date										
		Prescri	ption	Information							
Medication	Dose Strength ☐ 200mg/mL Vial Kit ☐ 200mg/	ml Startarkit		Direct	Qty	Refills					
	□ 200mg/mL prefilled Syringe	mi Stanter Kit		.oading Dose: Inject 400mg SC at Weeks 0, 2, and 4 faintenance Dose: Inject 200mg SC every 2 weeks							
Dupixent	□ PFS with needle shield 300 □ Prefilled Pen 300 mg/2 mL	mg/2 mL	□ Inject 300 mg Sub Q every week								
🗆 Entyvio	🗆 300mg vial		Loading Dose: Inject 300mg IV over 30 minutes at								
				Weeks 0, 2, and 6.							
				MaintenanceDose: Infuse 300mg IV over 30 minutes every 8 weeks							
🗆 Humira	Starter Kits:		Adult:								
	80mg/0.8mL Starter Pack Pre-Filled Pen			pading Dose: Inject 160mg	SC on Day 1, then 80mg						
	(Citrate Free) Maintenance:			on Day 15 (two weeks later) Maintenance Dose: Inject 40mg SC every other week							
	□ 40mg/0.4mL Pre-Filled Pen (Citrate Free) □ 40mg/0.4mL Pre-Filled Syringe (Citrate Free) □ Other:		(starting Day 29) Pediatric (>6 years and adolescents) 17kg to < 40kg □ Loading Dose: Inject 80mg SC on Day 1, 40mg on Day 15 (two weeks later)								
			MaintenanceDose:Inject20mgSCeveryotherweek (starting Day 29)								
								Pediatric (>6 years and adolescents) > 40kg			
			Loading Dose: Inject 160mg SC on Day 1, 80mg on Day 15 (two weeks later)								
							□ Maintenance Dose: Inject 40mg SC every other week				
				ting Day 29)							
Prescriber Signature:DAW (Dispense as Written) 🛛 Y 🖓 N Date:											

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately by calling or faxing back to the originator.

Crohn's and Ulcerative Colitis Referral Form		SUPERIOR BIOLOGICS Fax Referral To: 914-747-1170 Phone: 855-747-1150			SUPERIOR					
Date:					BIOLO	GIC	-0			
Patient Information				Prescriber Information						
Please complete the following or send patient demographic sl										
	:		—	City State Zin:						
				Phone:						
City, State, Zip				Fax:						
Home Phone:				DEA:NPI#:						
Cell Phone:										
DOB:										
Drive en de seu				nformation		0				
	ance:			#:						
				#:						
	Card:ID#:					Group:				
	ical Information (Section mus ation Insurance Number:	t be completed to	to proc	ess prescription)		(Attach separate sheet if	neede	ed)		
	Please include diagnosis name	with ICD-10 co	ode	Therapy Details	s : □ Ne	ew □ Reauthorization □ Re	estart			
 K50.00 Crohn's disease of small intestines without complicati K50.8 Crohn's disease of both intestines without complication K50.10 Crohn's disease of large intestines without complication K50.00 Crohn's disease, unspecified, without complication Other diagnosis: ICD-10 code 			ons ons 1s	Allergies Lab Data Prior Therapies		/lbs Height				
	Date of Description			Concomitant Medications Additional Comments						
Has a TB test been performed?						ed? □Yes □No				
	ent have an active infection?			njoolon naming	Jitequir					
	Review Date									
			otion	Information						
Medication	Dose Strength				Direction	ons	Qty	Refills		
 □ Avsola □ Inflectra □ Remicade □ Renflexis 	☐ 100mg Vial		 Loading Dose: Infuse 5mg/kg at Weeks 0, 2, and 6 Maintenance Dose: Infuse 5mg/kg every 8 weeks 							
□ Rinvoq	 Induction Therapy – 45 mg tablet Maintenance Therapy – 15 mg or 30 mg tablets 		 ☐ Induction Therapy: 45 mg PO daily x 8 weeks. Maintenance Therapy: ☐ 15 mg PO daily ☐ 30 mg PO daily 							
🗆 Simponi	□ 100mg/mL SmartJect Auto □ 100mg/mL Prefilled Syringe				-					
□ Stelara	☐ 130mg/26mL solution single ☐ 90mg/mL Prefilled Syringe Date of Initial Infusion:									
□ Skyrizi	use vial. Ongoing Therapy – 360 mg	vial. hor Ongoing Therapy – 360 mg/2.4 mL			I Initiation Therapy – □ Inject 600 mg IV over at least 1 Ir at Weeks 0, 4, 8. 1vial/week. I Ongoing Therapy – Week 12 – Inject 360 mg SC and ery 8 weeks thereafter. 1 device with prefilled cartridge.					
□ Xeljanz	 5mg tablet 10mg tablet 11mgXRtablet 22mgXRtablet 			aintenance Dose	∷ XR Nonce c	ng twice daily 🗆 XR: 22mg				
Prescriber Signature:DAW (Dispense as Written) 🛛 Y 🖓 N Date:										

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately by calling or faxing back to the originator.