

Ocrevus Referral) R U P	+ 20 (7 (& + \$ ' 9 \$ 1 & (' 7 + (5 \$ 3 , (6) D [5 H I H U U D O 7 R 3 K R Q H	
Date: _____		

3 \$ 7 , (1 7 , 1) 2 5 0 \$ 7 , 2 1	3 5 (6 & 5 , % (5 , 1) 2 5 0 \$ 7 , 2 1
Patient Name: _____	Prescriber Name: _____
Address: _____	Address: _____
City, State, Zip: _____	City, State, Zip: _____
Home Phone: _____	Phone: _____
Cell Phone: _____	Fax: _____
Date of Birth: _____	DEA#: _____ NPI#: _____
Gender: <input type="checkbox"/> M <input type="checkbox"/> F	Contact Person: _____

, 1 6 8 5 \$ 1 & (, 1) 2 5 0 \$ 7 (**Please attach the front and back of insurance and prescription drug card**)

Primary Insurance: _____	ID#: _____	Group: _____
Secondary Insurance: _____	ID#: _____	Group: _____
Prescription Card: _____	ID#: _____	BIN: _____ PCN: _____ Group: _____

, \$ * 1 2 6 , 6 , & '

G35 Relapsing forms of Multiple Sclerosis (Clinically isolated syndrome/relapsing-remitting disease/active secondary progressive disease)

G35 Primary Progressive Multiple Sclerosis

3 5 (6 & 5 ((1 , 1 *'

Hepatitis B Surface Antigen: _____

Total Hepatitis B Core Antibody (Anti-HBc): _____

Serum Immunoglobulins: _____ **AC OF CO EA T D**

Vaccination: _____ Height: _____ Weight: _____

Allergies: _____

(live or live-attenuated 4 weeks before, non-live 2 weeks before initiation of therapy)

Labs (During Therapy): _____

Vaccinations: Live-attenuated or live vaccines is not recommended during treatment and after discontinuation until B-cell repletion. Administer all necessary immunizations according to immunization guidelines at least 4 weeks prior to initiation for live or attenuated vaccines and at least 2 weeks prior to initiation for non-live vaccines.

Pre-screening: Required Hepatitis screening before first dose to include:

_____ Hepatitis B Surface Antigen (HBsAg) and Total Hepatitis B Core Antibody (anti-HBc) * Ocrevus® is contraindicated in patients with active HBV. Patients who are negative for surface antigen HBsAg (-) and positive for HB core antibody HBcAB (+) or positive for surface antigen HBsAg (+), should consult liver disease experts before starting and during treatment.

_____ Quantitative Serum Immunoglobulin Screening (IgG, IgA, IgM)

PRESCRIPTION ORDERS

<p>Premeds</p> <p>3 U H P H G L F O D W I B C 30 minutes prior to infusion:</p> <p><input type="checkbox"/> Acetaminophen PO: <input type="checkbox"/> 325mg <input type="checkbox"/> 500mg <input type="checkbox"/> 650mg</p> <p>Diphenhydramine: <input type="checkbox"/> 25mg IVP <input type="checkbox"/> 50mg IVP <input type="checkbox"/> 25mg PO <input type="checkbox"/> 50mg PO</p> <p>OR <input type="checkbox"/> Alternate oral antihistamine: <input type="checkbox"/> Cetirizine 10mg <input type="checkbox"/> Loratadine 10mg</p> <p><input type="checkbox"/> Fexofenadine 60mgs <input type="checkbox"/> Fexofenadine 180mgs</p> <p>IV Access Flush Order: NaCl 0.9% 5-10ml IV before and after infusion</p> <p><input type="checkbox"/> Methylprednisolone <input type="checkbox"/> 125mg IVP <input type="checkbox"/> 40mg IVP OR <input type="checkbox"/> _____mg PO</p> <p><input type="checkbox"/> Others/Miscellaneous: _____</p>	<p>Anaphylaxis Orders and Medications</p> <p>Diphenhydramine Administer 25 mg slow IV/IM may repeat x1 Dispense: 1 x 50 mg vial</p> <p>Epinephrine Autoinjector <input type="checkbox"/> Administer 0.15mg (1:2000) IM (< 30 Kg) <input type="checkbox"/> Administer 0.3mg (1:1000) IM (≥ 30 Kg)</p> <p>' L V S H Q V H: 1 package (2 pens)</p> <p>Sodium Chloride 0.9% <i>Use to maintain IV line, prevent or treat hypotension in case of anaphylaxis</i></p> <p>Dispense: QS</p>
---	---

0 H G L F D W L R Q

Ocrevus (Ocrelizumab) IV as directed to infuse per protocol via pump with 0.22 μ [: A E C A filter, following each infusion with a one hour post observation period.

Induction/Initial dosing: Induction/Initial dosing: 300mg Ocrevus IV in 250ml Sodium Chloride 0.9% to be infused at Week 0 over 2.5 hours or longer and 2 weeks later over 2.5 hours or longer. No Refills. ****To be infused in MD office or an Infusion suite.**

Maintenance dosing: 600mg Ocrevus IV in 500ml Sodium Chloride 0.9% to be infused every 6 months. 2 hrs or longer for eligible patients who have not experienced a serious infusion reaction with any previous Ocrevus Infusion 3.5-4 hrs or longer. Refills: X1 year ****Infusions to be performed under the close supervision of a healthcare professional and to observe the patient for least one hour after completion of the infusion.**

P { ^ A } ~ • q } A ~ a A T Ö A ~ a

By signing below, I certify that above therapy is medically necessary. 3 U H V F U L E H U ¶ V 6 L J Q D W X U H 6 , * 1 % (

Dispense as Written	Date	Substitution Allowed	Date
---------------------	------	----------------------	------

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately.